

Immunizations

Vaccines and Teens

The Busy Social Years

First Edition, 2011



The Children's Hospital of Philadelphia®



VACCINE EDUCATION CENTER

AMA
AMERICAN
MEDICAL
ASSOCIATION



**Teens have
their own
set of
health needs,**

You never thought it would get more hectic than it was when you were changing diapers, arranging play dates and helping with homework, but somehow it has. Adolescents and teens in the home bring a whole new definition to the word “busy.” They have their own schedules, deadlines and obligations. They also have their own set of health needs, including immunizations.

As children move into adolescence, they become increasingly susceptible to meningococcus, a disease that can claim a life in hours. They will soon begin exploring their sexuality, thereby increasing the likelihood of exposure to sexually-transmitted diseases, such as human papillomavirus. And their immunity from some childhood vaccines, such as pertussis (whooping cough) starts to wane. As a parent, how do you know what they need and when? Is it worth arguing over getting “a shot” and adding yet another appointment to the calendar? And, most importantly, are these vaccines safe?

In this booklet, we will discuss which vaccines are recommended, what diseases they prevent, and whether they are safe. We will also provide information about making your teen comfortable during immunizations and locating vaccine records for colleges and employers. With a better understanding of the unique needs of adolescents and teens, you will see why your doctor feels so strongly about the value of vaccines.

including immunizations.



Table of Contents

6	Why Do Teens Need Vaccines?
8	Which Vaccines Do Teens Need?
10	Vaccines Recommended for Teens <ul style="list-style-type: none">• Meningococcal Vaccine• Human Papillomavirus (HPV) Vaccine• Tetanus-Diphtheria-Acellular Pertussis (Tdap) Vaccine• Influenza Vaccine
18	A Word About Catch-up Vaccines <ul style="list-style-type: none">• Hepatitis A• Hepatitis B• Polio• Measles, Mumps and Rubella (MMR)• Varicella (Chickenpox)
21	Vaccines for Special Situations <ul style="list-style-type: none">• Vaccines for Special Work Environments• Vaccines for Teens with Chronic Diseases• Vaccines for Behaviors that Increase Risk• Vaccines Before Sexual Exploration• Vaccines for Sleepaway Camps, College and Traveling Abroad
23	Comforting Your Teen or Young Adult
24	Common Questions and Concerns About Vaccines
27	Vaccine Schedule
28	We Need What? Locating Vaccine Records
30	I Need More Information: Resources for Parents and Teens <ul style="list-style-type: none">• Vaccine-Related Information• Travel Information• Family Groups• Adolescent and Teen Resources• Social Media
34	Immunization Record

Why Do Teens Need Vaccines?

By adolescence, a child's immune system has been introduced to many viruses and bacteria; in some cases through illness or infection; in others, through immunization. Either way, their immune system is equipped to recognize these "repeat offenders" through its immunological memory.

Immunological memory is maintained in cells known as memory T and B cells. Memory T cells trigger a cascade of immune reactions that speed up the response to an infection, and memory B cells make antibodies to fight specific infections. Working together, these responses either prevent the infection completely or alter its course, leading to an illness that is less severe or of shorter duration.

Invincible Adolescents and Teens

Unfortunately (and contrary to their belief!) teenagers are not invincible. Their immune systems are not prepared to fight every infection. In fact, this age group is particularly susceptible to certain infections. Here's why:

- Fading immunity – Despite being immunized against diphtheria, tetanus and pertussis as young children, older children (and adults) become susceptible again. Unlike most diseases, protection provided by these vaccines is not lifelong.
- No previous exposure – While most vaccine-preventable diseases are spread casually through coughs, sneezes or shared objects, some, such as human papillomavirus (HPV), are only spread through intimate contact. Vaccines work best when they are introduced before an individual's first encounter with the virus, so that protective immunity is in place before exposure.



- Social habits – Adolescents and teens are inherently social. Unfortunately, the new experiences they seek — sleepaway camp, college dorms and night clubs — are often the same ones that increase their risk of infections with bacteria such as meningococcus.



- Changing viruses – Like children older than 6 months and most adults, adolescents and teens should get a yearly influenza vaccine. Because the virus changes so rapidly from season to season, the vaccine from the previous year may not provide immunity.

Sending Them Out Into the World

As parents, we want to prepare our kids for whatever they may encounter when we're not around. We give them our values and discuss good decision-making. We teach them how to cross the street safely and to use bike helmets and seat belts. We monitor their friends, grades and eating habits. Vaccines provide one more opportunity for protection.

While all vaccines are made to afford protection without causing illness, various approaches are used to make vaccines against different types of bacteria and viruses:

- Use a piece of the virus or bacteria – HPV and meningococcal vaccines are made this way. The meningococcal vaccine is linked to another protein that helps make the immune response stronger.
- Use the toxins produced by the bacteria – The bacteria that cause diphtheria, tetanus and whooping cough all produce toxins that cause disease. Therefore, in order to be protective, the vaccines contain inactivated forms of the toxins, known as toxoids.
- Use the whole, killed virus – The influenza shot employs this method.
- Use the live, weakened virus – The intranasal version of the influenza vaccine employs this method.

Which Vaccines Do Teens Need?

Around 2005, several new vaccines became available that made immunization of older children more common:

- **Meningococcus:** The meningococcus vaccine was recommended for all 11- to 12-year-olds.

Emerging data suggests that immunity with the newer version of the meningococcal vaccine is not as long-lasting as originally expected; therefore, teens who received the vaccine at 11 to 12 years of age should get a second dose at 16 to 18 years of age. Thirteen- to 15-year-olds who get the vaccine also need a booster dose between 16 and 18 years of age, and 16- to 18-year-olds who have not had the vaccine before should get one dose.

- **Tdap:** The existing tetanus-diphtheria (Td) vaccine, recommended every 10 years, was modified to include a pertussis or whooping cough component (Tdap) that will protect adolescents and adults from three diseases instead of two. This change was important not only for the recipients of the vaccine who could suffer months of coughing spells and broken ribs from whooping cough, but also for infants who were not old enough to be protected from suffering and death caused by pertussis.

Adolescents between 11 and 12 years of age should get a single dose of Tdap vaccine. Teens between 13 and 18 years of age who have not had the vaccine should also get one dose.

Older children and adults were not recommended to receive the infant version of the vaccine, known as DTaP, because 2 or 3 of every 100 recipients experienced significant swelling of the arm or leg in which the vaccine was given. The newer version contains quantities of diphtheria and pertussis that make this side effect much less likely. The lower quantities in the formulation are indicated by the lower case “d” and “p” in the name: Tdap.

- **Human papillomavirus:** An HPV vaccine was licensed in 2006. HPV is the most common sexually transmitted disease in the United States and the world. HPV is a known cause of cervical and anal cancer and genital warts as well as occasional cancers of the head and neck. Because the vaccine only works *before* someone is infected and because 11- to 12-year olds are already recommended to get other vaccines, the HPV vaccine was recommended for this age group as well.

HPV vaccine requires three doses. The second dose is given 1 to 2 months after the first dose. The third dose is given 6 months after the first dose.

Children as young as 9 and young adults up to 26 can receive this vaccine.

- **Influenza:** Recommendations have gradually expanded to include more people who should receive influenza vaccine every year. By 2009, children 6 months to 18 years of age were recommended to receive the influenza vaccine, and in 2010, the recommendation expanded to include everyone over 6 months of age.

- First entrance into middle school, high school or college is a great time to catch up on vaccines. Check to make sure your child is immune to hepatitis A, hepatitis B, polio, measles, mumps, rubella and chickenpox.

Originally the HPV vaccine was only recommended for girls, but studies have now shown it is also safe and effective in boys. Giving the vaccine to boys protects them against genital warts and some cancers. Also, transmission of the virus will be decreased between sexual partners.



Vaccines Recommended for Teens

Meningococcal Vaccine

What is meningococcus?

Meningococcus is a bacterium that lives on the lining of the nose and throat of about 1 in 10 people. Most people don't have any symptoms of infection.

What is my child's risk of getting meningococcus?

Before the current meningococcal vaccine, every year approximately 2,500 people in the United States got meningococcus and about 120 died from their infections. Approximately 400 people who survived were left with permanent disabilities, such as seizures, loss of limbs, kidney disease, deafness and mental retardation.

Although children younger than 2 years old are the most likely to catch

People who have recently had a disturbance to the lining of their throat and those who are in close contact with a large group of people are at increased risk of infection with meningococcus. Several groups fall into these categories, including people who had a recent infection of the nose, throat or lungs; college freshmen living in dorms; military recruits living in barracks; people exposed to tobacco smoke or indoor wood stoves; people who go to bars or nightclubs; and people who share drinking glasses or cigarettes.

meningococcus, teens are the most likely to die from it.

How do you catch meningococcus?

Meningococcal bacteria are passed from person to person by coughing or sneezing.

What is the meningococcal vaccine?

The meningococcal vaccine is made using purified sugars from the protein coating the bacteria and from four of the five different types of meningococcus. The vaccine also includes a harmless protein that isn't part of meningococcus but helps improve immunity to the meningococcal sugars in the vaccine. Because the protein is linked to the sugars, it's called the meningococcal conjugate vaccine.

Does the meningococcal vaccine prevent all cases of meningitis?

No. Meningitis refers to an infection of the lining of the brain and spinal cord. However, meningococcal bacteria are not the only cause of meningitis, so while preventing meningococcal infections will reduce the number of cases of meningitis, it will not prevent all cases of meningitis. While a few causes of meningitis are prevented by vaccines given in infancy, such as *Haemophilus influenzae* type b and pneumococcus, others are not vaccine-preventable.

Further, the existing meningococcal vaccine does not protect against all types of meningococcal bacteria.



Is the meningococcal vaccine safe?

Yes, the meningococcal vaccine is safe:

- Some teens may develop minor side effects such as pain or redness at the site of injection, as well as fever, headache or tiredness.
- Since some teens are prone to fainting following receipt of vaccines, they should stay seated or lie down at the doctor's office for about fifteen minutes after getting a vaccine.
- Some have questioned whether the meningococcal vaccine causes Guillain-Barré Syndrome (GBS), a disease involving muscle weakness, burning or tingling of the limbs, loss of muscle tone, and paralysis. Studies have shown that the meningococcal vaccine does not cause GBS.

Do the benefits of the meningococcal vaccine outweigh its risks?

Every year, people in the United States are severely debilitated or die from infections with meningococcal bacteria. Since the vaccine does not cause severe reactions, the benefits clearly outweigh the risks.



Human Papillomavirus (HPV) Vaccine

What is human papillomavirus?

Human papillomavirus (HPV) infects the genital area and the lining of the cervix. There are many types of HPV, and while most people do not even realize they were infected, in some cases, infections last for a long time. Some types of HPV cause genital warts, and other types cause cancers of the cervix, head, neck, anus, vagina, vulva and penis. Often the cancers do not develop until 20 to 25 years after the infection.

What is my child's risk of getting HPV?

Before the HPV vaccine, every year in the United States 6 million people were newly infected with HPV. About half of these infections occurred in girls and young women between 15 and 24 years of age. Because about 40 of 100 girls are infected with HPV in their first two years of sexual activity, it is important to get the HPV vaccine before the start of sexual exploration.

About 20 million people in the United States are already infected. About 10,000 women are diagnosed with cervical cancer each year, and 4,000 die.

How do you catch HPV?

HPV is transmitted by genital contact, most often, but not only, during sex. Although condoms are helpful, they are not completely protective.

Because about 40 of 100 girls are infected with HPV in their first two years of sexual activity, it is important to get the HPV vaccine before the start of sexual exploration.

What is the HPV vaccine?

The HPV vaccine is made using a protein from the surface of HPV. Two HPV vaccines are available; both contain the protein from two types of HPV that most commonly cause cervical cancer, and one contains proteins from two additional types of HPV that commonly cause genital warts. Girls can get either version; however, boys are only recommended to get the version that protects against four types of HPV.

Because initial exposure to HPV does not typically include all four types in the vaccine, young people who have begun sexual exploration may still benefit from receiving this vaccine.

Does the HPV vaccine work?

Because HPV does not typically cause cervical cancer until decades after infection, people have wondered how scientists know the vaccine works. In fact, the answer can be found in what is seen when cervical cells are examined under the microscope following Pap smears. When women are infected with HPV and the infection does not clear quickly, cells of the cervix begin to develop precancerous and then cancerous changes, called cervical intraepithelial neoplasia (CIN). The HPV vaccine has been shown to prevent CIN, a required step in the development of cervical cancer.

Is the HPV vaccine safe?

Yes. Side effects following receipt of the HPV vaccine can include pain, swelling or redness at the injection site. Some teenagers faint after receiving the vaccine.

Some people have attributed blood clots, strokes, heart attacks, chronic fatigue syndrome and even deaths to the HPV vaccine. However, studies have shown the vaccine did not cause any of these conditions.

Do girls and women who get the HPV vaccine still need to get Pap tests?

Yes. Because HPV vaccine does not protect against all types of HPV that cause cervical cancer, women can still be infected and later develop this disease, so they should continue to use this important preventive measure.

Do those who get HPV vaccine still need to worry about sexually transmitted diseases?

Yes. HPV vaccine does not protect against all types of HPV, nor does it prevent other types of sexually transmitted infections such as syphilis, gonorrhea, chlamydia and herpes.

Do the benefits of the HPV vaccine outweigh its risks?

Every year, thousands of people are infected with HPV, and some will die from their infections. Since the vaccine does not cause any serious reactions, the benefits of the vaccine clearly outweigh the risks.

Tetanus-Diphtheria-Acellular Pertussis (Tdap) Vaccine

Tetanus

What is tetanus?

Tetanus is a bacterium that typically infects older adults. Tetanus makes a toxin that causes severe and painful spasms of the muscles, including muscles of the jaw. For this reason, tetanus is sometimes referred to as lockjaw. Muscle spasms of the throat can block the windpipe and cause instant death from suffocation. The tetanus toxin can also cause severe and permanent damage to the heart. About three of every 10 people who get tetanus die from the disease.

What is my child's risk of getting tetanus?

Each year about 40 people in the United States get tetanus and several die.

How do you get tetanus?

Tetanus bacteria live in the soil and can enter the skin after a cut or puncture wound. Because tetanus bacteria will always be present in the soil, the risk of getting infected will never go away. Further, because the disease is not passed from person to person, it doesn't matter how many other people in the community are immunized, an individual's risk of getting tetanus will remain the same.

Diphtheria

What is diphtheria?

Diphtheria is caused by a bacterium that typically infects children and adolescents. However, recent outbreaks in other countries have occurred primarily among adults due to fading immunity.

The diphtheria bacterium causes a thick coating on the back of the throat that makes it difficult to swallow and breathe. Also, the bacterium makes a harmful protein, called a toxin, that can invade the heart, kidneys and nervous system.

What is my child's risk of getting diphtheria?

About 2 to 5 people in the United States get diphtheria each year.

How do you catch diphtheria?

Diphtheria is very contagious and is spread by coughing and sneezing.

Pertussis

What is pertussis?

Pertussis is a type of bacteria that infects people of all ages. The bacterium makes several toxins that cause a thick, sticky mucus that clogs the windpipe and causes painful coughing spasms. When a person breathes in against the narrowed windpipe, it causes a whooping sound. This is why pertussis is often called whooping cough. In young infants this struggle to breathe can be deadly. In adults coughing spasms

can lead to broken ribs, seizures and hernias. Pertussis can also cause pneumonia, seizures and permanent brain damage.

What is my child's risk of getting pertussis?

Pertussis is quite common, but often is under-diagnosed. Estimates are that about 800,000 cases occur each year in the United States, with about 15 to 20 deaths.

How do you catch pertussis?

Pertussis is spread through coughing and direct contact with respiratory secretions.

Tdap Vaccine

What is the Tdap vaccine?

Tdap vaccine protects against three different bacteria. The vaccine is made by taking the toxins produced by each of these bacteria and inactivating them with a chemical. The chemical is then washed away, leaving the purified, inactivated toxins (called toxoids).

Tdap vaccine is different from Td vaccine because it includes the whooping cough component. Tdap

vaccine is different from DTaP vaccine given to infants and young children because it contains lesser quantities of diphtheria and pertussis vaccines.

Does the Tdap vaccine prevent pertussis?

Yes. In medical studies, Tdap vaccine was shown to prevent pertussis in 9 of every 10 recipients.

Is Tdap vaccine safe?

A small number of people who receive Tdap vaccine will experience pain, redness or swelling at the injection site, as well as slight fever, headache, fatigue or upset stomach. Less often, and particularly in adults, an exaggerated local reaction occurs that may involve swelling from the shoulder to the elbow.

Do the benefits of the Tdap vaccine outweigh its risks?

Every year in the United States, people are infected with these three diseases, and some die from them. Although side effects sometimes occur, these reactions do not result in permanent harm or death, so the benefits of the vaccine outweigh the risks.



Influenza Vaccine

What is influenza?

Influenza is a virus that infects the respiratory system, attacking the lining of the nose, windpipe, large breathing tubes, small breathing tubes and lungs.

What is my child's risk of getting influenza?

Each year in the United States approximately 200,000 people are hospitalized and thousands to tens of thousands die from influenza. Unfortunately, some of those who are hospitalized and die are previously healthy children and teens.

How do you catch influenza?

Influenza is transmitted by coughing and sneezing as well as by direct contact with infected respiratory secretions.



What is the influenza vaccine?

There are two different influenza vaccines:

- The inactivated influenza vaccine (or flu shot) is made by taking influenza viruses, growing them in eggs, purifying them and chemically inactivating them so they cannot cause disease.
- The live, weakened influenza vaccine (FluMist®) is made so it cannot grow in the lungs. However, because the weakened influenza viruses can grow in the lining of the nose, they cause an excellent protective immune response. FluMist is also made by growing the viruses in eggs. This vaccine is given as a nasal spray.

Both vaccines are remade annually so that they contain the three different strains of influenza viruses expected to cause disease that year.

Does the influenza vaccine work?

Yes. Because the strains of influenza in the vaccine are based on those circulating in other parts of the world, some years the vaccine is a better match (and more effective) than others. However, the influenza vaccine typically prevents about 70 to 85 of every 100 recipients from developing a severe influenza infection.

Is the influenza vaccine safe?

People who get the influenza shot may experience pain, redness and swelling at the injection site. The nasal spray vaccine may cause cough, runny nose, congestion or sore throat in a small number of recipients. In rare cases, people experience an allergic reaction such as hives, most likely caused by egg proteins in the vaccine.

Although the influenza vaccine is made in eggs and some people are severely allergic to eggs, the quantity of egg proteins in the vaccine is insufficient to cause a severe allergic response. But just to be sure, people with severe egg allergies should remain at the provider's office for 30 minutes after receiving the influenza vaccine.

Do the benefits of the influenza vaccine outweigh its risks?

Influenza virus kills thousands to tens of thousands of people every year, some of whom were previously healthy. In addition, about 200,000 people are hospitalized each year with fever, croup, bronchitis, bronchiolitis or pneumonia caused by influenza. Because the vaccine does not cause serious reactions, the benefits of the vaccine clearly outweigh the risks.

Because other viruses cause respiratory illnesses and colds, it is important to remember the influenza vaccine will only protect against influenza viruses.

A Word About Catch-up Vaccines

Hepatitis A

Hepatitis A is a virus that infects the liver. While most people who get it don't have any symptoms, for some infection with hepatitis A causes loss of appetite, vomiting, nausea, fatigue and jaundice (yellowing of the eyes and skin). In rare cases, hepatitis A causes an overwhelming infection of the liver resulting in death. This virus can spread either from one person to another or through contaminated food or water.

The hepatitis A vaccine is made by purifying hepatitis A virus and inactivating it with a chemical. The vaccine, now recommended for all children at age 1, is given as a shot in two doses separated by six months.

Side effects are minor and can include a headache or pain, warmth or swelling at the injection site.

Hepatitis B

Hepatitis B is a virus that infects the liver. Many people who get hepatitis B do not know they have it, but some have long-term, silent infections that lead to cirrhosis or liver cancer. Symptoms of hepatitis B infection include loss of appetite, vomiting, nausea, fatigue and jaundice. Hepatitis B can spread through sexual contact, from infected mothers to their babies during birth, and by sharing items that may contain small quantities of blood or saliva, such as toothbrushes, razors or nail clippers.

The hepatitis B vaccine is made from the protein coat that surrounds the virus, called the hepatitis B surface protein. The vaccine is given as a shot in two or three doses. Because the vaccine was recommended at birth beginning in 1992, many adolescents and teens have already had it; however, those who have not should be caught up. In most cases, three doses of hepatitis B vaccine will be needed with the second dose being given 1 to 2 months after the first and the third dose, 4 months after the first. Some 11- to 15-year olds may receive a two-dose version in which the second dose is given 4 to 6 months after the first.

Side effects are typically minor and include pain, redness and tenderness at the injection site, fever, headache, fatigue and irritability. Rarely, recipients have a severe allergic reaction with hives, rash or a drop in blood pressure. This type of reaction only occurs in about 1 of every one million recipients and usually within 30 minutes of getting the vaccine.

Polio

Most adolescents and teens have received the polio vaccine, but if they did not receive the recommended number of doses, they need to catch up. Polio is caused by a virus that in most cases causes a mild intestinal infection. However, in about 1 of every 200 people the virus also infects the brain and spinal cord, leading to paralysis. Polio is spread when

contaminated food, hands or objects are put in the mouth; it is quite contagious.

The polio vaccine is made from the three different types of polio that infect people. These viruses are purified and inactivated with a chemical so that they cannot cause disease. The vaccine is given as a shot. Adolescents and teens not previously immunized need three doses. Those who received some doses previously may require fewer doses.

The vaccine may cause pain, redness or tenderness at the site of the injection.

Measles, Mumps and Rubella (MMR)

Measles, mumps and rubella are three viruses that cause distinct illnesses. Measles is extremely contagious and can be transmitted through viral particles that remain in the air as long as two hours after an infected person leaves. Mumps and rubella are both transmitted through respiratory secretions such as coughs and sneezes. All three infections are extremely dangerous for pregnant women;

therefore, it is important for women to be immunized before they decide to start having children.

Measles, mumps and rubella vaccines are made from weakened, live viruses that can no longer cause disease. Vaccine viruses are combined to form the MMR vaccine. MMR vaccine is given as a shot and requires two doses. In adolescents and teens the doses are separated by 4 weeks. Since some in this age group already had one dose at a younger age, they may only require a single, second dose.

The MMR vaccine may cause mild side effects such as pain, redness or tenderness at the site of injection as well as fever or rash in some recipients. About 1 of every 24,000 people who get the vaccine may have a temporary decrease in platelets, which are cells in the bloodstream that help blood to clot. Temporary arthritis or swelling of the joints may also occur in a small number of vaccine recipients, mostly adolescent girls and women.



Varicella (Chickenpox)

Varicella, also known as chickenpox, is a virus that causes fever and about 300 to 500 blisters over the entire body. In severe cases the virus can also cause pneumonia, encephalitis (swelling of the brain) or skin infections from “flesh-eating” bacteria. Teens and adults tend to experience more complications. Infection during pregnancy can lead to birth defects or spontaneous abortion and fetal death. Chickenpox is spread by coughing and sneezing and is quite contagious.

The chickenpox vaccine is made from a live, weakened form of the virus. The vaccine is given as a shot in two doses. Because the recommendation for a second dose is relatively new, some adolescents and teens will have had only one dose as a younger child, therefore, requiring one additional dose.

The vaccine causes pain, redness or tenderness at the site of injection, and a small number of recipients will develop a few blisters.



Vaccines for Special Situations

As teens have new opportunities and develop into young adults, new encounters may present health risks.

Vaccines for Special Work Environments

If your child is employed in a healthcare setting or in a position in which he or she may come into contact with blood, it is important to be up-to-date on all vaccines, particularly those that prevent diseases transmitted by blood or body fluids from an infected person, such as hepatitis B.

If your child is employed in a setting in which he or she is coming into contact with large groups of people, he or she should be up-to-date on vaccines that prevent highly contagious diseases such as pertussis, measles, influenza and chickenpox.

If your child is employed in an outdoor setting, such as lawn care, farming or landscaping, he or she should have recently received a tetanus-containing vaccine since tetanus bacteria live in the soil and can cause an infection after entering the body through a wound.

Vaccines for Teens with Chronic Diseases

Unfortunately, even young people are sometimes diagnosed with illnesses that compromise their immune systems or require them to take medications that can make them more susceptible to vaccine-preventable diseases. For example, people with asthma are more susceptible to

diseases that infect the lungs, such as influenza and pneumococcus.

If your child is being treated for diseases such as diabetes, asthma, chronic lung disease, kidney disease, heart disease, sickle cell disease, cancer, or immune diseases such as HIV or AIDS, you should check with your healthcare provider to determine whether your teenager needs any vaccines, and whether there are vaccines that they shouldn't get during their treatment. For example, some people treated with steroids should not get live, weakened viral vaccines such as chickenpox or MMR.

Vaccines for Behaviors That Increase Risk

Because teens often believe they are invincible, they do not always take care of themselves. When they get run down and fail to eat a well-balanced diet, they may be more susceptible to certain illnesses. In addition, activities such as smoking, sharing drinks, getting tattoos or piercings, sharing personal care items (such as toothbrushes, razors or nail clippers), or sharing illegal drug paraphernalia can make young adults more likely to catch vaccine-preventable diseases such as pneumococcus, meningococcus and hepatitis B.

Vaccines Before Sexual Exploration

Although parents do not like to think about teens and their sexual exploration, the reality is that they will be involved in sexual encounters eventually. Parents can make sure their children are protected from certain situations by having them vaccinated before these encounters begin:

- Girls should have received rubella (the “R” in MMR) and chickenpox vaccines before they become pregnant because if infected with these diseases during pregnancy, unborn babies can be harmed or die.
- If a teenager is pregnant during influenza season, she should have the influenza shot since she will be more susceptible to complications of influenza if infected.
- Both boys and girls should receive human papillomavirus (HPV) and hepatitis B vaccines. HPV and hepatitis B can be spread through sexual contact.

Vaccines for Sleepaway Camps, College and Traveling Abroad

When your children begin adventures that take them away from home, it is possible they will come in contact with viruses and bacteria to which they have not been exposed. For example, because meningococcal bacteria live harmlessly in the noses and throats of about 1 of every 10 people, your child may be exposed to a type that he or she is susceptible to when sharing living quarters with others.

If your child has the opportunity to travel abroad, he or she may require additional vaccines or medications. The best way to determine this is to take your child to a travel medicine clinic a few months before the trip. Factors such as where they are going, where they will stay, and what activities they will participate in will determine their health needs. You can find more information about travel resources, including how to find a travel clinic near you, in the resources section of this booklet (page 30).



Studies have shown that college freshmen living in dorms are five times more likely to get a meningococcal infection compared with peers not living on campus.

Comforting Your Teen or Young Adult

As difficult as it might have been to watch your infant and young child get vaccines, taking your older child or teen for vaccines can be worse if they are needle-phobic. You can no longer hold them on your lap and comfort them — even getting them to the appointment may be difficult. However, here are some things you may try:

For You:

- As with other issues, if your child sees that vaccines are important to you, they will be more accepting of them. Before the appointment, discuss getting vaccines with your child, answer their questions, and reassure them.
- During the appointment, bring the immunization record, read materials provided by the office staff, ask questions and check about the use of pain medication following the appointment.
- After the appointment, provide lots of fluids; realize that your child's appetite may decrease over the next 24 hours; use a cool, wet cloth on a red or swollen injection site; have your child take a cool bath or shower for fever; provide pain medication as instructed by your doctor; and call the office if an unusual reaction occurs.

For Your Child:

- Take something or someone to the appointment that will serve as a distraction. Talking to a parent, sibling or friend, listening to music, texting or playing games on a cell phone, or reading a book may take your child's mind off of the impending shot(s).
- Tell your child to try to relax his muscles and not to look while the needle is being given. Taking a few short, deep breaths and then a few longer breaths during the procedure will also help.
- Since the sensation of cold usually overpowers that of pain, ask for an alcohol pad, have your child rub it on her wrist, and have her blow on the spot while the shot is given.
- Finally, remind your child that the pain of a needle is nothing compared with the diseases that vaccines prevent.

Adolescents and teens must be comfortable getting vaccines without the use of physical constraint.

Common Questions and Concerns About Vaccines

Are vaccines safe?

To best answer this question, we must first define what we mean when we say “safe.” If by “safe” we mean completely risk-free, then vaccines aren’t 100 percent safe. Like all medicines, vaccines have mild side effects, such as pain, tenderness or redness at the site of injection. And some vaccines have very rare, but more serious, side effects.

But nothing is harmless. Anything that we put into our bodies (like vitamins or antibiotics) can have side effects. Even the most routine activities can be associated with hidden dangers. For example, consider seat belts. It’s possible that in an accident a seat belt could cause a minor injury, like a bruise. But if you measure the risk of wearing a seat belt against the risk of not wearing one, the decision to wear a seat belt is an easy one. Similarly, for each of the recommended vaccines, the benefits far outweigh the risks.

Do we still need vaccines?

Yes. Vaccines are still given for three reasons:

- For common diseases (like chickenpox or pertussis), a choice not to get a vaccine is a choice to risk natural infection. For example, every year hundreds of thousands of people are infected with pertussis and some die from the disease. Therefore, it’s important to get the vaccine.
- Some diseases (like measles or mumps) still occur in the United




States at low levels. If immunization rates drop further, these diseases will come roaring back.

- While some diseases (like polio, rubella or diphtheria) have been either completely or virtually eliminated from the United States, they still occur in other parts of the world. Polio still commonly paralyzes people in Africa, and rubella still causes birth defects and miscarriages in many parts of the world. Because international travel is common, these diseases are only a plane ride away from coming back into the United States.

Why do adolescents and teens need vaccines?

Older children and teens need vaccines for several reasons:

- To boost immunity – Some vaccines do not provide enough immunity to last a lifetime, so additional doses are necessary as children (and adults) get older. The Tdap vaccine is one example.
- To protect against diseases that have not been encountered – Some vaccines protect against viruses the person hasn't seen before. The HPV vaccine is an example.
- To protect against viruses that change – Some viruses adapt to their environment in a “survival of the fittest” manner, so that older versions of the vaccine are no longer effective in protecting against the disease. Influenza vaccine is the only example.
- To protect against diseases that tend to infect particular age groups – For example, the meningococcal vaccine is important because the chance of getting this disease decreases after age 2, but begins to increase again around adolescence.



Do vaccines weaken the immune system?

No, in fact, vaccines prevent infections that weaken the immune system. Because bacteria and viruses contained in vaccines are highly weakened versions of natural bacteria and viruses, they do not weaken the immune system.

On the contrary, infections with natural viruses can weaken the immune system. For example, people infected with influenza virus are at risk of developing severe bacterial pneumonia. Also, people infected with chickenpox virus are at risk of developing severe infections of the skin caused by “flesh-eating” bacteria.

Can vaccines cause long-term diseases like multiple sclerosis, diabetes or asthma?

Studies have shown that vaccines don't cause autism, diabetes, multiple sclerosis, allergies, asthma or permanent brain damage.

When one event precedes another, we often wonder whether they are related. For example, some people who smoke a lot of cigarettes get lung cancer. But when scientists wanted to know whether cigarette smoking caused lung cancer, several studies performed in the 1950s and 1960s compared the likelihood of lung cancer in people who smoked cigarettes with that in people who didn't smoke. The best studies matched these two groups of people with regard to age, general health, medications and so on. By matching these groups, they made sure the only difference between them was cigarette smoking. The result was clear: Cigarette smoking caused lung cancer.

Similarly, some people who use cell phones get brain cancer. To answer the question of whether cell phones caused brain cancer, the occurrence of brain cancer in people who used cell phones was compared with that in people who didn't. Again, these groups were matched to make sure the only difference between them was cell phone use. The result was also clear: Cell phones didn't cause brain cancer.

Because vaccines are given to nearly everyone, many people with chronic diseases will have received vaccines. And some of these people will have received vaccines close in time to the appearance of the chronic condition. The question is: "Did the vaccine cause the disease?" The best way to answer this question is to do studies similar to those described for smoking and cell phones. Although not all potential associations have been studied, many have; therefore, we know that vaccines don't cause autism, diabetes, multiple sclerosis, allergies, asthma or permanent brain damage.



Vaccine Schedule

Vaccine	Ages 11-12	Ages 13-18
Meningococcus	1st dose at 11-12yo* 2nd dose at 16-18yo	13-15yo: 1 dose, followed by a booster at 16-18yo 16-18yo: 1 dose
Tdap	1 dose	1 dose if not previously received
HPV	2nd dose 1-2 months after first 3rd dose 6 months after first	3 doses if not previously received Remaining doses if immunization was started

Ages 11-18

Influenza	1 dose annually
Hepatitis A†	2 doses
Hepatitis B†	2 or 3 doses
Polio†	3 doses
Measles, mumps, rubella (MMR)†	1 or 2 doses
Chickenpox†	1 or 2 doses

*yo=years old

† Catch-up shots may be required depending on the child's vaccination history.



We Need What? Locating Vaccine Records

As your child's world expands to include sleepaway camps, first jobs and college, you may find that people are asking for their vaccine record.

While some parents have a copy of their child's immunization record, many do not. So what should you do if you can't find your child's record?

- Check with your child's most recent healthcare provider. If your child has switched doctors, you may need to check with previous providers, especially those who cared for your child before he or she entered school.
- Check with your child's school since they sometimes keep copies.
- If your state health department has an immunization registry, you may be able to get a copy from them.
- Check your child's baby books and your old health records as you may have a copy of the record in a place you hadn't thought about.
- Old school or camp forms may also have the information.

What if I can't find my child's immunization record?

If you are unable to locate your child's immunization record, he or she may need to:

- Have a blood test – A blood test can help determine immunity to some vaccine-preventable diseases, such as measles, mumps, rubella, chickenpox, hepatitis A, hepatitis B, tetanus, diphtheria and polio. However, if the blood test reveals that the teenager is not immune to one or more of these diseases, he or she will need to be vaccinated. Since an extra dose of vaccine will not cause harm, some parents opt to get the vaccines in question without getting the blood test.
- Be vaccinated – Some vaccines may be necessary either because there is not a blood test to determine immunity or because the blood test reveals vaccines that are needed.

While many parents are concerned that getting vaccines will be harmful if their child already had been vaccinated, they can be reassured that extra doses of vaccine, like an exposure to the disease, will serve to strengthen the immune system.



There may be costs associated with the blood test and extra co-pays for visits to make up vaccines. Each vaccine has its own set of requirements regarding how soon after the initial dose subsequent doses can be administered.

Once you and the healthcare provider decide how to proceed, you can set up a schedule to be sure the vaccine series is completed. In addition to the records the doctor keeps, ask them to complete a record that you take each time you go for additional doses.

I Need More Information: Resources for Parents and Teens

Vaccine-related Information

Internet Resources

- Centers for Disease Control and Prevention (CDC)
www.cdc.gov/vaccines/who/teens
- Vaccine Education Center (VEC) at The Children’s Hospital of Philadelphia
www.vaccine.chop.edu
- American Medical Association (AMA)
www.ama-assn.org/go/roadmaps-adolescents
- Vaccine Information for the Public and Health Professionals (a program of Immunization Action Coalition – IAC)
www.vaccineinformation.org
- National Network for Immunization Information (NNii)
www.immunizationinfo.org
- Vaccinate Your Baby (a program of Every Child by Two – ECBT)
www.vaccinateyourbaby.org

Books

- *Vaccines and Your Child: Separating Fact from Fiction* (Columbia University Press, 2011) co-written by Paul A. Offit, M.D., and Charlotte A. Moser



- *The Complete Idiot's Guide to Vaccinations* (Penguin Group USA, 2009) co-written by Michael J. Smith, M.D., M.S.C.E., and Laurie Bouck

- *Do Vaccines Cause That?! A Guide for Evaluating Vaccine Safety Concerns* (i4ph, 2008) co-written by Martin G. Meyers, M.D., and Diego Pineda

Pamphlets and Information Sheets

- Vaccine Information Statements (VIS) on all vaccines are offered by the CDC.

www.cdc.gov/vaccines/pubs/vis

- Various informational tear sheets are available for download from the Vaccine Education Center at The Children's Hospital of Philadelphia.

www.vaccine.chop.edu/resources



Travel Information

- CDC Travel Website – Includes information specific to destinations, vaccines and diseases

www.cdc.gov/travel

- Find a travel clinic – Both the International Society of Travel Medicine (ISTM) and the American Society of Tropical Medicine and Hygiene (ASTMH) provide clinic locators on their websites:

ISTM: www.istm.org

ASTMH: www.astmh.org

Family Groups

- Parents PACK (Possessing, Accessing and Communicating Knowledge about Vaccines) – A program directed by the Vaccine Education Center at The Children's Hospital of Philadelphia. Offers a free monthly email newsletter about vaccines. Vaccine information is provided for all age groups.

www.vaccine.chop.edu/parents

- Vaccinate Your Baby – An immunization awareness campaign administered by Every Child By Two (ECBT); includes information about vaccines and their safety.

www.vaccinateyourbaby.org

- Families Fighting Flu – A nonprofit public organization composed of families and pediatricians who have experienced firsthand what it is like to lose a child to flu or to have a child experience severe medical complications from flu.

www.familiesfightingflu.org

- Meningitis Angels – A nonprofit organization composed of families affected by meningitis and dedicated to the support of other victims of bacterial meningitis and their families.

www.meningitis-angels.org

- National Meningitis Association – A nonprofit organization founded by parents whose children have died or live with permanent disabilities from meningococcal disease.

www.nmaus.org

- PKIDS – A national nonprofit organization that offers support to families whose children are living with hepatitis, HIV/AIDS, or other chronic, viral infectious diseases and that educates the public about infectious diseases.

www.pkids.org

- Tamika & Friends – A national nonprofit organization dedicated to raising awareness about cervical cancer and its links to human papillomavirus.

www.tamikaandfriends.org

- Pearl of Wisdom Campaign – A united, global effort to raise awareness of the opportunities now available to prevent cervical cancer.

www.pearlofwisdom.us

- The Yellow Umbrella Organization – An organization that “is about working together, under the same umbrella, toward a common goal ... It’s about cervical cancer prevention.”

www.theyellowumbrella.org



Adolescent and Teen Resources

- Getvaxed.org – An initiative of PKIDS that provides immunization-related information and videos for teens and young adults.

www.getvaxed.org

- Pirls – A project of PKIDS, Pirls is a website for girls where they can get health information, play games, win prizes and listen to Pirls radio. Pirls is also on Facebook, Twitter and MySpace.

www.pirls.pkids.org

- Stomping Out Meningitis – An initiative of Meningitis Angels that brings meningitis survivors, teen family members



and friends together to share personal stories and educate other teens about the importance of preventing meningococcal meningitis through healthy living and vaccination.

www.stompingoutmeningitis.com

Social Media

- Shot of Prevention – a blog hosted by ECBT. Go to *shotofprevention.com*. Also on Facebook and Twitter.
- Parents Who Protect – a blog hosted by the National Meningitis Association. Go to *parentswhoprotect.com*
- PKIDS is on Facebook, Twitter, Flickr, YouTube, Delicious and other social media sites. They also have a blog. Learn more at *www.pkids.org*.
- The Pearl of Wisdom Campaign and the Yellow Umbrella Organization are on Facebook, YouTube and Twitter.
- Families Fighting Flu and the National Meningitis Association are on Facebook and Twitter.

Immunization Record

Below is a list of the recommended vaccines for adolescents and teens. Please have your child's physician or other healthcare professional fill in the date that your child receives a vaccine.

Child's name _____ Date of birth _____

Vaccine	Number of Doses		
Tdap			
Meningococcus			
Human Papillomavirus			
Influenza			

Are there any vaccines to catch up on?

Check that your child had the appropriate number of doses of the following vaccines by having your healthcare provider add the information from your child's chart here:

Hepatitis A _____

Hepatitis B _____

Polio _____

MMR _____

Varicella _____





Vaccines and Teens: The Busy Social Years was written and produced by the Vaccine Education Center at The Children's Hospital of Philadelphia in collaboration with the American Medical Association. The Center was formed in October 2000 to provide accurate, comprehensive and up-to-date information about vaccines and the diseases they prevent. The Vaccine Education Center is funded by endowed chairs from The Children's Hospital of Philadelphia. The center does not receive support from vaccine companies.

For more information about vaccines,
visit the Vaccine Education Center websites at
vaccine.chop.edu
and
vaccine.chop.edu/parents

 The Children's Hospital of Philadelphia®
Hope lives here.

34th Street and Civic Center Boulevard, Philadelphia, PA 19104-4399
1-800 TRY CHOP www.chop.edu

Founded in 1855, The Children's Hospital of Philadelphia is the birthplace of pediatric medicine in America. Throughout its history, a passionate spirit of innovation has driven this renowned institution to pursue scientific discovery, establish the highest standards of patient care and train future leaders in pediatrics. For a century and a half, Children's Hospital has served as a haven of hope for children and families worldwide.

The Children's Hospital of Philadelphia and the  logo are registered marks of The Children's Hospital of Philadelphia.

Keep the connection.

